

References

- Aaronson, J., Abramovitz, S., Smiley, R., Tangel, V., & Landau, R. (2017). A survey of intravenous remifentanyl use for labor analgesia at academic medical centers in the United States. *Obstetric Anesthesiology*, *124*(4), 1208–1210. <https://doi.org/10.1213/ANE.0000000000001622>
- Ansari, J., Carvelho, B., Shafer, S., & Flood, P. (2016). Pharmacokinetics and pharmacodynamics of drugs commonly used in pregnancy and parturition. *Anesthesia & Analgesia*, *122*(3), 786-804. <https://doi.org/10.1213/ANE.0000000000001143>
- Balki, M., Kasodekar, S., Dhumne, S., Bernstein, P., & Carvalho, J. C. (2007). Remifentanyl patient-controlled analgesia for labour: Optimizing drug delivery regimens. *Canadian Journal of Anaesthesia*, *54*(8), 626-633. <https://doi.org/10.1007/BF03022956>
- Balcioglu, O., Akin, S., Demir, S., & Aribogan, A. (2007). Patient-controlled intravenous analgesia with remifentanyl in nulliparous subjects in labor. *Expert Opinion on Pharmacotherapy*, *8*(18), 3089-3096. <https://doi.org/10.1517/14656566.8.18.3089>
- Brennan, F., Obs, D., Carr, D.B., & Cousins, M. (2007). Pain management: a fundamental human right. *Anesthesia & Analgesia*, *105*(1), 205-221. <https://doi.org/10.1213/01.ane.0000268145.52345.55>
- Butwick, A. J., Wong, C. A., & Guo, N. (2018). Maternal body mass index and use of labor neuraxial analgesia. *Anesthesiology*, *129*(3), 448–458. <https://doi.org/10.1097/aln.0000000000002322>
- Centers for Disease Control and Prevention. (2021, March 2). *FastStats - Births and Natality*. Centers for Disease Control and Prevention. <https://www.cdc.gov/nchs/fastats/births.htm>

- Dershwitz, M., Randal, G.I., Rosow, C.E., Fragen, R.J., Connors, P.M., Librojo, E.S., Shaw, D.L., Peng, A.W., & Jamerson, B.D. (1995). Initial clinical experience with remifentanyl, a new opioid metabolized by esterases. *Anesthesia Analgesia*, *81*(3), 619-623.
[https://doi.org/ 10.1097/00000539-199509000-00035](https://doi.org/10.1097/00000539-199509000-00035)
- Devabhakthuni, S. (2013). Efficacy and safety of remifentanyl as an alternative labor analgesic. *Clinical Medicine Insights: Women's Health*, *6*(2013). [https://doi.org/ 10.4137/CMWH.S8015](https://doi.org/10.4137/CMWH.S8015)
- D'Onofrio, P., Novelli, A. M. M., Mecacci, F., & Scarselli, G. (2009). The efficacy and safety of continuous intravenous administration of remifentanyl for birth pain relief: an open study of 205 parturients. *Anesthesia & Analgesia*. *109*(6), 1922 – 1924.
[https://doi.org/ 10.1213/ane.0b013e3181acc6fc](https://doi.org/10.1213/ane.0b013e3181acc6fc)
- Douma, M. R., Verwey, R. A., Kam-Endtz, C. E., Van der Linden, P. D., & Stienstra, R. (2010). Obstetric analgesia: A comparison of patient-controlled meperidine, remifentanyl, and fentanyl in labour. *British Journal of Anesthesia*, *104*(2), 209-215. [https://doi.org/ 10.1093/bja/aep359](https://doi.org/10.1093/bja/aep359)
- Evron, S., Glezerman, M., Sadan, O., Boaz, M., & Ezri, T. (2005). Remifentanyl: A novel systemic analgesic for labor pain. *Anesthesia & Analgesia*. *100*(1), 233 – 238.
[https://doi.org/ 10.1213/01.ANE.0000143351.64538.BC](https://doi.org/10.1213/01.ANE.0000143351.64538.BC)
- Glass, P.S., Gan, T.J., & Howell, S. (1999). A review of the pharmacokinetics and pharmacodynamics of remifentanyl. *Anesthesia & Analgesia*, *89*(4), 7. [https://doi.org/ 10.1097/00000539-199910001-00003](https://doi.org/10.1097/00000539-199910001-00003)
- Gupta, S. & Partani, S. (2018). Neuraxial techniques of labour analgesia. *Indian Journal of Anaesthesia*, *62*(9). 658-666. https://doi.org/10.4103/ija.IJA_445_18

- Ismail, M. T. & Hassanin, M. Z. (2012). Neuraxial analgesia versus intravenous remifentanil for pain relief in early labor in nulliparous women. *Archives of Gynecology and Obstetrics*, 286(6), 1375 – 1381. <https://doi.org/10.1007/s00404-012-2459-3>
- Jelting, Y., Weibel, S., Afshari, A., Pace, N. L., Jokinen, J., Artmann, T., Eberhart, L. H. J., & Kranke, P. (2017). Patient-controlled analgesia with remifentanil vs. alternative parental methods for pain management in labour: A Cochrane systemic review. *Anaesthesia*, 72(8), 1016–1028. <https://doi.org/10.1111/anae.13971>
- Jia, Z., Li, Y., Jia, H., Ren, J., & Xie, N. (2020). Curative effect of remifentanil on labor analgesia in newborns. *Journal of Maternal-Fetal & Neonatal Medicine*, 33(11), 1913-1918. <https://doi.org/10.1080/14767058.2018.1533946>
- Jost, A., Ban, B., & Karmenik, M. (2012). Modified patient-controlled remifentanil bolus delivery regimen for labour pain. *Association of Anaesthetists*, 68, 245-252. <https://doi.org/10.1111/anae.12038>
- Kan, R.E., Hughes, S.C., Rosen, M.A., Kessin, C., Preston, P.G., Lobo, E.P. (1998). Intravenous remifentanil: Placental transfer, maternal and neonatal effects. *Anesthesiology*, 88(6), 1467-1474. <https://doi.org/10.1097/00000542-199806000-00008>
- Kapila, A., Glass, P. S., Jacobs, J. R., Muir, K. T., Hermann, D. J., Shiraishi, M., Howell, S., & Smith, R. L. (1995). Measured context-sensitive half-times of remifentanil and alfentanil. *Anesthesiology*, 83(5), 968-975.
- Karadjova, D., Shosholcheva, M., Ivanov, E., Sivevski, A., Kjaev, I., Kartalov, A., Kuzmanovska, B., Spasova, R., Kocovski, G. & Aleksiovska-Papestiev, I. (2019). Side effects of intravenous patient-controlled analgesia with remifentanil compared with

- intermittent epidural bolus for labour analgesia: A randomized controlled trial. *PRILOZI*, 40(3), 99-108. <https://doi.org/10.2478/prilozi-2020-0009>
- Karol, D. & Weiniger, C. F. (2021). Update on non-neuraxial labor analgesia. *Current Anesthesiology Reports*, 11, 348 – 354. <https://doi.org/10.1007/s40140-021-00463-4>
- Konefal, H., Jaskot, B., Czeszynska, M. B., & Pastuszka, J. (2013). Remifentanil patient-controlled analgesia for labor-monitoring of newborn heart rate, blood pressure, and oxygen saturation during the first 24 hours after delivery. *Archives of Medical Science*, 9(4), 677 – 702.
- Lee, M., Zhu, F., Moodie, J., Zhang, Z., Cheng, D., & Martin, J. (2017). Remifentanil as an alternative to epidural analgesia for vaginal delivery: A meta-analysis of randomized trials. *Journal of Clinical Anesthesia*, 39(2017), 57–63. <https://doi.org/10.1016/j.jclinane.2017.03.026>
- Lehmann, K. A. (2005). Recent developments in patient-controlled analgesia. *Journal of Pain Symptom Management*, 29(5), 72-89.
- Leong, W. L., Leong Sng, B., & Heng Sia, A. T. (2011). A comparison between remifentanil and meperidine for labor analgesia: A systemic review. *Anesthesia & Analgesia*, 113(4), 818-825. <https://doi.org/10.1213/ANE.0b013e3182289fe9>
- Liu, Z.Q., Chen, X.B., Li, H.B., Qiu, M.T., & Duan, T. (2014). A comparison of remifentanil parturient-controlled intravenous analgesia with epidural analgesia: A meta-analysis of randomized controlled trials. *Anesthesia & Analgesia*, 118(3), 598 – 603. <https://doi.org/10.1213/ANE.0000000000000077>

- Lin, R., Tao, Y., Yu, Y., Xu, Z., Su, J., & Liu, Z. (2014). Intravenous remifentanil versus epidural ropivacaine sufentanil for labour analgesia: A Retrospective Study. *PLoS ONE*, 9(11). <https://doi.org/10.1371/journal.pone.0112283>
- Logtenberg, S. L. M., Vink, M. L., Godfried, M. B., Beenackers, I. C. M., Schellevis, F. G., Mol, B. W., & Verhoeven, C. J. (2019). Serious adverse events attributed to remifentanil patient-controlled analgesia during labour in the Netherlands. *International Journal of Obstetric Anesthesia*, 39, 22-39. <https://doi.org/10.1016/j.ijoa.2018.10.013>
- LoMauro, A. & Aliverti, A. (2015). Respiratory physiology of pregnancy. *Breathe*, 11, 197-301. <https://doi.org/10.1183/20734735.008615>
- Markley, J. C. & Rollins, M. D. (2017). Non-neuraxial labor analgesia: options. *Clinical Obstetrics & Gynecology*, 60 (2), 350-364. <https://doi.org/10.1097/GRF.0000000000000277>
- Marwah, R., Hassan, S., Carvalho, J. C., & Balki, M. (2012). Remifentanil versus fentanyl for intravenous patient-controlled labour analgesia: an observational study. *Canadian Journal of Anesthesia*, 59, 246-254. <https://doi.org/10.1007/s12630-011-9625-0>
- Melber, A. A. (2018a). Remifentanil patient-controlled analgesia (PCA) in labour – in the eye of the storm. *Association of Anaesthetists*. 74, 277-279. <https://doi.org/10.1111/anae.14536>
- Melber, A. A., Jelting, Y., Huber, M., Keller, D., Dullenkopf, A., Girard, T., & Kranke, P. (2018b). Remifentanil patient-controlled analgesia in labour: six-year audit of outcome data of RemiPCA SAFE Network (2010 – 2015). *International Journal of Obstetric Anesthesia*. 39, 12-21. <https://doi.org/10.1016/j.ijoa.2018.12.004>

- Messmer, A. A., Potts, J. M., & Orlikowski, C. E. (2016). A prospective observation study of maternal oxygenation during remifentanyl patient-controlled analgesia use in labour. *Anaesthesia*, *71*, 171-176. <https://doi.org/10.1111/anae.13329>
- Murray, H., Hodgkinson, D., Hughes, D. (2019). Remifentanyl patient-controlled intravenous analgesia during labour: A retrospective observational study of 10 years' experience. *International Journal of Obstetric Anesthesia*, *39*, 29-34. <https://doi.org/10.1016/j.ijoa.2019.05.012>
- Ohashi, Y., Baghirzada, L., Sumikura, H., & Balki, M. (2016). Remifentanyl for labor analgesia: A comprehensive review. *Japanese Society of Anesthesia*. *30*, 1020-1030. <https://doi.org/10.1007/s00540-016-2233-y>
- Osterman, M., & Martin, J.A. (2011). Epidural and spinal anesthesia use during labor: 27-state reporting area, 2008. *National vital statistics reports: from the Centers for Disease Control and Prevention, National Center for Health Statistics, National Vital Statistics System*. *59* (5), 1-13.
- Pillay, M., Shaw, M., Vincent, C., & Frawley, N. (2019). Remifentanyl in labour: Retrospective audit of a single regional centre's outcomes. *Journal of Paediatrics and Child Health*, *55*, 3 – 55.
- Raja, S.N., Carr, D.B., Cohen, M., Finnerup, N.B., Flor, H., Gibson, S., Keefe, F.J., Mogil, J.S., Ringkamp, M., Sluka, K.A., Song, X., Stevens, B., Sullivan, M.D., Tutelman, P.R., Ushida, T., & Vader, K. (2020). The revised International Association for the Study of Pain definition of pain: concepts, challenges, and compromises. *Pain*, *161*(9), 1976-1982. <https://doi.org/10.1097/j.pain.0000000000001939>

- Schnol, H., Paul, N., & Belfer, I. (2014). Labor pain mechanisms. *International Anesthesiology Clinic*, 52(3), 1 – 17. [https://doi.org/ 10.1097/AIA.0000000000000019](https://doi.org/10.1097/AIA.0000000000000019)
- Shen, M. K., Wu, Z. F., Zhu, A. B, Shen, X. F., Yang, J. J., & Feng, S. W. (2013). Remifentanil for labour analgesia: A double-blinded, randomized controlled trial of maternal and neonatal effects of patient-controlled analgesia versus continuous infusion. *Anaesthesia*, 68(3), 236 -244. [https://doi.org/ 10.1111/anae.12098](https://doi.org/10.1111/anae.12098).
- Soens, M. A., Birback, D. J., Ranasinghe, J. S., & Van Zundert, A. (2008). Obstetric anesthesia for the obese and morbidly obese patient: an ounce of prevention is worth more than a pound of treatment. *Acta Anaesthesiology Scandinavia*, 52, 6 – 19. [https://doi.org/ 10.1111/j.1399-6576.2007.01483.x](https://doi.org/10.1111/j.1399-6576.2007.01483.x)
- Stocki, D., Matot, I., Einav, S., Eventov-Friedman, S., Ginosar, Y., & Weiniger, C.F. (2014). A randomized controlled trial of the efficacy and respiratory effects of patient-controlled intravenous remifentanil analgesia and patient-controlled epidural analgesia in laboring women. *Anesth Analg*, 118(3), 589-597. <https://doi.org/10.1213/ANE.0b013e3182a7cd1b>
- Stourac, P., Suchomelova, H., Stodulkova, M., Huser, M., Krikava, I., Janku, P., Haklova, O., Hakl, L., Stoudek, R., Gal, R., & Sevcik, P. (2014). Comparison of parturient-controlled remifentanil with epidural bupivacaine and sufentanil for labor analgesia: A randomized controlled trial. *Biomedical Papers of the Medical Faculty of the University of Palacky, Olomouc, Czech Republic*, 158(2), 227 – 232.
- Sugur, T., Kizilates, E., Kizilates, A., Inanoglu, K & Karsli, B. (2020). Labor analgesia: Comparison of patient-controlled analgesia and intravenous patient-controlled analgesia. *Agri*, 32(1), 8 – 18. [https://doi.org/ 10.14744/agri.2019.35403](https://doi.org/10.14744/agri.2019.35403)

- Tveit, T. O., Seiler, S., Halvorsen, A., & Rosland, J. H. (2012). Labour analgesia: A randomized, controlled trial comparing intravenous remifentanyl and epidural analgesia with ropivacaine and fentanyl. *European Journal of Anaesthesiology*, 29, 129-136. <https://doi.org/10.1097/EJA.0b013e32834dfa98>
- United Regional. (2018). *Remifentanyl PCA Protocol for Labor Patient*.
- Van de Velde, M. & Carvalho, B. (2016). Remifentanyl for labor analgesia: an evidence-based narrative review. *International Journal of Obstetric Anesthesia*, 25, 66-74. <https://doi.org/10.1016/j.ijoa.2015.12.004>
- Varposhti, M. R., Ahmadi, N., Masoodifar, M., Shahshahan, Z., & Tabatabaie, M. H. (2013). Comparison of remifentanyl: Entonox with Entonox alone in labor analgesia. *Advanced Biomedical Research*, 2, 648-655. <https://doi.org/10.4103/2277-9175.122511>
- Volmanen, P., Akural, E., Raudaskoski, T., Ohtonen, P., & Alahuhta, S. (2005). Comparison of remifentanyl and nitrous oxide in labour analgesia. *Anaesthesiologica Scandinavica*, 49(4), 453-458. <https://doi.org/10.1111/j.1399-6576.2005.00639.x>
- Welzing, L., Ebenfeld, S. E., Dlugay, V., Wiessen, M. H. J., Roth, B., & Mueller, C. (2011). Remifentanyl degradation in umbilical cord blood of preterm infants. *Anesthesiology*, 114(3), 570-577. <https://doi.org/10.1097/ALN.0b013e318204e043>
- Wilson, M. J. A., MacArther, C., Hewitt, C. A., Handley, K., Gao, F., Beeson, L., & Daniels, J. (2018). Intravenous remifentanyl patient-controlled analgesia versus intramuscular pethidine for pain relief in labour (RESPITE): an open-label, multicentre, randomized controlled trial. *Lancet*. 392, 662-672. [http://doi.org/10.1016/S0140-6736\(18\)31613-1](http://doi.org/10.1016/S0140-6736(18)31613-1)

Zhang, P., Yu, Z., Zhai, M., Cui, J., & Wang, J. (2021). Effect and safety of remifentanyl patient-controlled analgesia compared with epidural analgesia in labor: An updated meta-analysis of randomized controlled trials. *Gynecologic and Obstetric Investigation, 86*, 231-238. [https://doi.org/ 10.1159/000515531](https://doi.org/10.1159/000515531)